

## INSTRUCTIONS FOR COLONOSCOPY: MIRALAX/GATORADE

**DATE:** \_\_\_\_\_ **TIME:** \_\_\_\_\_

**PLEASE REPORT TO:** \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

**\*Arrive 45 minutes prior to procedure time to register\***

**PLEASE MAKE ARRANGEMENTS TO HAVE SOMEONE DRIVE YOU HOME BECAUSE YOU WILL BE MEDICATED FOR THIS PROCEDURE.** If you are using public transportation, a responsible person must accompany you to your home. If these instructions are not followed, your procedure will be cancelled.

If you have a history of heart disease and take a daily **aspirin**, you **do not** need to stop taking it. If however, you have no history of heart disease and are taking a daily aspirin for preventative purposes, please stop this one week before your procedure. Be sure to check with our office regarding any of the following: diabetes, valvular heart disease, bleeding tendencies or anti-coagulation therapy.

**ANTI-INFLAMMATORY MEDICATIONS** such as Advil, Motrin, Aleve, Naprosyn, and other Ibuprofen products, as well as **IRON PILLS**, and **METAMUCIL/FIBER SUBSTITUTES** should be stopped **ONE-WEEK** prior to your procedure. (Calcium, Tylenol, Vitamins C & B are OK to continue taking).

Two days prior to your procedure, avoid foods such as fruits, vegetables, skins, roughage, seeds and nuts.

You may have **CLEAR LIQUIDS ONLY** for one full day before your procedure. (This means **NO SOLID FOOD** from the time you wake up in the morning the day before the procedure). Clear liquids include broth, decaf coffee/tea (no cream, sugar is OK), clear juices (like apple or white grape), all sodas, Jello, Gatorade, Popsicles, and water. **NO RED DYES:** Jello, Popsicles, sherbert, and juices must be non-red in color, as red dyes found in foods can look like blood during the procedure.

**AT 3:00PM** the day before your procedure take **two (2) Dulcolax (Bisacodyl) tablets**. These can be purchased at the pharmacy without a prescription.

**AT 5:00PM** the day before your procedure mix the 255 gram bottle of **MiraLax powder** (you will be given a prescription for this) in a 64oz bottle of Gatorade. Shake the solution until the MiraLax is dissolved. Drink an 8oz glass of this mixture every 15 minutes until the solution is gone.

You may continue with **clear liquids until 4 hours prior to your scheduled procedure time** (for example, if your procedure is at 12:00pm, you may have clear liquids until 8:00am). **After this time, DO NOT HAVE ANYTHING BY MOUTH.** Do not take any medications the morning of the procedure unless otherwise instructed by our office.

If you have any questions or difficulties with this preparation, please contact our office at 401-421-8800.

Most insurance companies cover this procedure however if you have not checked with your insurance company or if you are unsure of your benefits please check with your carrier regarding coverage.